

ON THE MAXIMUM OF A STATIONARY INDEPENDENT INCREMENT PROCESS

SHELDON M. ROSS

AD 734133



OPERATIONS RESEARCH CENTER



COLLEGE OF ENGINEERING
UNIVERSITY OF CALIFORNIA • BERKELEY

NATIONAL TECHNICAL: INFORMATION SERVICE Springfield, Va. 22151

ON THE MAXIMUM OF A STATIONARY INDEPENDENT INCREMENT PROCESS

Ъу

Sheldon M. Ross
Department of Industrial Engineering
and Operations Research
University of California, Berkeley



NOVEMBER 1971

ORC 71-29

This research has been partially supported by the Office of Naval Research under Contract N00014-69-A-0200-1010 and the U.S. Army Research Office-Durham under Contract DA-31-124-ARO-D-331 with the University of California. Reproduction in whole or in part is permitted for any purpose of the United States Government.

Security Classification											
DOCUMENT CONT											
Security classification of title, body of abstract and indexing a 1. Official ting Activity (Corporate author)	annotation must be e		uverall report in classified) ECURITY CLASSIFICATION								
			ssified								
University of California, Berkeley		26. GROUP	9911164								
3 REPORT TITLE											
ON THE MAXIMUM OF A STATIONARY INDEPENDENT	r increment	PROCESS									
	,	. 110 0===									
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)											
Research Report											
5 AUTHOR(5) (First name, middle initial, last name)											
Sheldon M. Ross											
Sherdon m. Ross											
6 REPORT DATE	74. TOTAL NO. OF	E BAGES	76. NO. OF REFS								
November 1971	6	FRULL	3								
NOVEMBER 1971 BA. CONTRACT OR GRANT NO.	98. ORIGINATOR'S	S REPORT NUMI									
DA-31-124-ARO-D-331		ORC 71-29	•								
6. PROJECT NO.		OKC /1-2,	9								
20014501B14C		- 141 /Any o	to and land								
с.	this report)	₹T NO(3) (Any	ther numbers that may be assigned								
d.											
10. DISTRIBUTION STATEMENT											
This document has been approved for publi	to volesse s	ا بواده ا-	to distribution is								
unlimited.	.C lereast	id bare,	is distribution to								
13. SUPPLEMENTARY NOTES Also supported by the	12. SPONSORING M	TARY ACTI									
Office of Naval Research under Contract			Office-Durham								
N00014-69-A-0200-1010.	Box CM, Du	uke Station	n								
	Durham, No	orth Caroli	ina 27706								
13 ABSTRACT											
SEE ABSTRACT.											
SEE ABSTRACT.											
			3								
		i									

DD FORM .. 1473 (PAGE 1)

Unclassified
Security Classification

S/N 0101-807-6811

A-3140A

Security Classification		LINKA		. LINK B		LINK	
KEY WORDS		ROLE		ROLE	WT	HOLE	-
Stationary Independent Increment I	rocess						l
Renewal Process							
					!		
Ladder Height					ļ		
				1			
·							
				'			
							!
					1		
	٠.					1	
]				
					1		
			ł				
					1		
				ı			
		[

DD FORM 1473 (BACK)

Unclassified
Security Classification

ABSTRACT

A stationary independent increment process is the continuous time analogue of the discrete random walk, and, as such, has a wide variety of applications. In this paper we consider M(t), the maximum value that such a process attains by time t. By using renewal theoretic methods we obtain results about M(t). In particular we show that if μ , the mean drift of the process, is positive, then $\frac{M(t)}{t}$ converges to μ , and $E\big(M(t+h)-M(t)\big)+h\mu$.

ON THE MAXIMUM OF A STATIONARY INDEPENDENT INCREMENT PROCESS

by

Sheldon M. Ross

Let $\{X(t)$, $t\geq 0\}$ be a separable stationary independent increment process continuous in probability and with X(0)=0. Suppose that $E[X(t)]=\mu t$ where $\mu>0$, and let

$$M(t) = \sup_{0 \le s \le t} X(s) .$$

Theorem 1:

- (i) $M(t)/t^{a+s}$, μ as $t \rightarrow \infty$
- (ii) $EM(t)/t + \mu$ as $t + \infty$.

Proof:

Fix some constant b>0 , and define T_n , J_n , $n\geq 1$, by

$$T_1 = \inf \{t : X(t) \ge b\}$$
, $X(T_1) = b + J_1$
 $T_n = \inf \{t : X(t) \ge b + X(T_{n-1})\}$, $X(T_n) = b + X(T_{n-1}) + J_n$, $n > 1$.

Now, due to the stationary independent increment assumption, it follows that $\{T_n-T_{n-1}\ ,\ n\geq 1\} \ \text{is a renewal process. Letting } \ N(t) \ \text{be the number of renewals by time } t \ \text{for this process, i.e., } \ N(t) = \max \left\{n: T_n \leq t\right\} \ , \text{ we have } t \in \mathbb{R}^n \times \mathbb{R}^n \times$

(1)
$$bN(t) + J_1 + ... + J_{N(t)} \le M(t) \le bN(t) + J_1 + ... + J_{N(t)} + b$$
.

Now, from the strong law of large numbers and the renewal result $N(t)/t \stackrel{a_{\downarrow}s}{\longrightarrow} 1/ET_1$, we have that

(2)
$$1/t (bN(t) + J_1 + ... + J_{N(t)})^{a_1 s_1} \cdot \frac{b + EJ_1}{ET_1}$$

Now, from a similar theorem in discrete time (i.e., Stein's lemma), it follows that $ET_1 < \infty$ and thus from Martingale theory

(3)
$$E[b + J_1] = E[X(T_1)] = \mu ET_1$$
,

and hence (i) follows from (1), (2) and (3). To prove (ii), we first note that

$$1/t E[bN(t) + J_1 + ... + J_{N(t)}] = bEN(t)/t + \frac{EJ_1(EN(t) + 1)}{t} - \frac{E[J_{N(t)+1}]}{t}$$

by Wald's equation. However by regarding J_i as a reward earned at the time of the ith renewal it follows, since $EJ_1 < \infty$, by standard arguments ([1], Page 53) that $\frac{E[J_N(t)+1]}{t} \to 0$. Hence, by the elementary renewal theorem,

$$1/t E[bN(t) + J_1 + ... + J_{N(t)}] + \frac{b + EJ_1}{ET_1} = \mu$$

and (ii) is proven.

Q.E.D.

Theorem 2:

If $E[X^2(t)] < \infty$, then, for each h > 0,

$$E[M(t + h) - M(t)] \rightarrow h\mu .$$

Proof:

By (1) we have

$$b(N(t+h) - N(t)) + J_1 + \dots + J_{N(t+h)} - (J_1 + \dots + J_{N(t)}) - b \le M(t+h) - M(t) \le b(N(t+h) - N(t)) + J_1 + \dots + J_{N(t+h)} - (J_1 + \dots + J_{N(t)}) + b.$$

Taking expectations, and using Wald's equation, yields

$$(b + EJ_1)E[N(t + h) - N(t)] + E[J_{N(t)+1}] - E[J_{N(t+h)+1}] - b \le$$

$$E[M(t + h) - M(t)] \le (b + EJ_1)E[N(t + h) - N(t)] + E[J_{N(t)+1}] -$$

$$E[J_{N(t+h)+1}] + b .$$

We shall show in the ensuing lemma that $\lim_{t\to\infty} E[J_{N(t)+1}]$ exists and is finite. Hence, the result follows from the above by first applying Blackwell's theorem and then letting $b\to 0$.

Q.E.D.

Lemma 1:

If $E[X^2(t)] < \infty$, then

$$\lim_{t\to\infty} E[J_{N(t)+1}] = \frac{E[J_1T_1]}{E[T_1]} < \infty .$$

Proof:

Let $g(t) = E[J_{N(t)+1}]$ and let F denote the distribution of T_1 . Then

$$g(t) = \int_{0}^{\infty} E[J_{N(t)+1} \mid T_{1} = x]dF(x)$$

$$= \int_{0}^{t} g(t - x)dF(x) + \int_{t}^{\infty} E[J_{1} \mid T_{1} = x]dF(x) .$$

This renewal type equation has the solution

$$g(t) = h(t) + \int_0^t h(t - x) dm(x)$$

where

$$h(t) = \int_{t}^{\infty} E[J_1 \mid T_1 = x]dF(x)$$
.

From the key renewal theorem it follows that

$$\lim_{t \to \infty} g(t) = \frac{\int_{0}^{\infty} h(t)dt}{E[T_{1}]} = \frac{\int_{0}^{\infty} \int_{t}^{\infty} E[J_{1} \mid T_{1} = x]dF(x)dt}{E[T_{1}]} = \frac{\int_{0}^{\infty} xE[J_{1} \mid T_{1} = x]dF(x)}{E[T_{1}]} = \frac{E[J_{1}T_{1}]}{E[T_{1}]}$$

where the interchange is justified by the nonnegativity of rewards. That $\mathrm{E}[\mathrm{J}_1\mathrm{T}_1] < \infty$ follows from a similar result in discrete time. (The discrete time result states that for a random walk with positive drift whose steps have finite variance, the variance of the value of the process the first time it exceeds b is finite. The equivalent discrete time version of $\mathrm{E}[\mathrm{J}_1\mathrm{T}_1] < \infty$ then follows from Stein's lemma upon application of the Cauchy-Schwarz inequality.)

Q.E.D.

Remarks:

The technique employed in the above proof is similar to the ladder variable approach in fluctuation theory. That is, we say that a ladder point is reached not when the process attains a new high but when it passes the previous ladder height by at least some fixed value. It is to be hoped that this technique will prove useful in studying other properties of stationary independent increment processes. Also this approach should be contrasted with the approach, recently developed by M. Rubinovitch, in which a point t is said to be a ladder point if M(t) = X(t). By applying the theory of regenerative events, as developed by Kingman, Rubinovitch

is then able to obtain many interesting results.

For a discrete time proof of (i), the interested reader should consult Heyde.

人名英格兰斯 医克勒氏试验检尿病 医克勒氏病 医克勒氏病 医克勒氏病 医克勒氏病 医克勒氏病 医二种结合 医二种生物 医二种生物 医克格特氏病 医牙管丛皮肤

REFERENCES

- [1] Ross, S., APPLIED PROBABILITY MODELS WITH OPTIMIZATION APPLICATIONS, Holden-Day, (1970).
- [2] Heyde, C.C., "Some Renewal Theorems with Application to a First Passage Problem," Annals of Mathematical Statistics, Vol. 37, pp. 699-711, (1966).
- [3] Rubinovitch, M., "Ladder Regenerative Events with Applications to Dam Models," Technical Report No. 58, Department of Operations Research, Cornell University, (1968).